
## Readings of the radiation rate with the cooperation of universities

Upper column: Reading of the integrated dose(24h) Lower column: the reference value which was calculated as the number per one hour

Prefecture	Monitoring	City	4/10 ~ 4/11
Prefecture	Point	City	
Hokkaido	1	Muroran City	2 μ Sv (0.08 μ Sv/h)
	2	Obihiro City	1 μ Sv (0.04 μ Sv/h)
	3	Asahikawa City	1 μ Sv (0.04 μ Sv/h)
	4	Kitami City	1 μ Sv (0.04 μ Sv/h)
	5	Kushiro City	1 μ Sv (0.04 μ Sv/h)
	6	Hakodate City	1 μ Sv (0.04 μ Sv/h)
Aomori	7	Hirosaki City	1 μ Sv (0.04 μ Sv/h)
	8	Hachinohe City	Less than 1 μ Sv
Miyagi	9	Sendai City	3 μ Sv (0.13 μ Sv/h)
Yamagata	10	Yonezawa City	2 µ Sv (0.08 µ Sv/h)
	11	Tsuruoka City	1 μ Sv (0.04 μ Sv/h)
Fukushima	12	Fukushima City	10 µ Sv (0.42 µ Sv/h)
Ibaraki	13	Tsukuba City	3 µ Sv (0.13 µ Sv/h)
Tochigi	14	Oyama City	3 µ Sv (0.13 µ Sv/h)
Gunma	15	Kiryu City	3 μ Sv (0.13 μ Sv/h)
Chiba	16	Chiba City	3 μ Sv ( 0.13 μ Sv/h )
	17	Kisarazu City	3 μ Sv (0.13 μ Sv/h)
Tokyo	18	Bunkyo Ward	2 µ Sv (0.08 µ Sv/h)
	19	Fuchu City	2 µ Sv (0.08 µ Sv/h)
	20	Meguro Ward	2 µ Sv (0.08 µ Sv/h)
	21	Minato Ward	2 µ Sv (0.08 µ Sv/h)
	22	Hachioji City	2 µ Sv (0.08 µ Sv/h)
Kanagawa	23	Yokohama City	2 µ Sv (0.08 µ Sv/h)
Niigata	24	Nagaoka City	2 µ Sv (0.08 µ Sv/h)
Nagano	25	Matsumoto City	3 µ Sv (0.13 µ Sv/h)
	26	Ueda City	2 µ Sv (0.08 µ Sv/h)
-l- M. I	the integrated	dose(24h) from around 2PM to the next	

<sup>\*</sup> We have measured the integrated dose(24h) from around 2PM to the next day.

<sup>\*</sup>Readings of lower column are the reference value because of the lower limit of the pocket dosimeter (1  $\mu$  Sv)



% Sv [Sievert]=Constant of organism effect by kind of radiation(%)  $\times$  Gy [gray]

X It is 1 in case of X ray and  $\gamma$  ray.

MEXT makes this, based on "Nuclear power 2002" made by Agency of Natural Resources and Energy.