Vomit Clean-up Procedure



- Using the original solution may produce harmful gases, **so it must be diluted with water before use.** (Make sure to check a product label since every product has a different concentration)

form of original solution.

- Label the container to ensure that children do not accidentally drink it.

Vomit Clean-up Procedure

3

Clean-up Procedure

1

Dilute sodium hypochlorite with a specified amount of water.



Cover the vomit with paper towels and apply disinfectant.

2



6

Wipe towards the center to gather it.



4

Put the matter wiped in Step [3] in doubled plastic bags, and apply disinfectant.



5

Spray the disinfectant on paper towels to wipe the surrounding areas for disinfection.



Put all items used in the doubled plastic bag. Close each bag tightly for disposal.





- Open a window for ventilation.
- Do not let people enter the room for 30 minutes after the clean-up procedure.
- Wash your hands thoroughly after the clean-up procedure.

Issued on: 2025/01/24

Issued by: the Office for a Professional Platform of Health and Hygiene Environment in Schools for Foreign Students provided by MEXT (mediPhone Inc.) Please recognize that this document is prepared on January 24, 2025 by the Office for a Professional Platform of Health and Hygiene Environment in Schools for Foreign Students provided by MEXT, and this tentative translation may vary from a standard translation published by the Japanese Government.

